Closedness of the solution mapping for parametric equilibrium problems

M. Bogdan* and D. Inoan†

Abstract

The equilibrium problems are treated as classical inequations to obtain closedness of the solution map defined on the set of parameters.

AMS subject clasification: 49J93D

Key words: Equilibrium problems, Closed set-valued maps, Pseudomonotonicity.

1 Introduction

This work is motivated by two recent papers of Bogdan and Kolumbán [2] and Yu et al. [9]. The first one deals with pseudomonotone operators defined on Sobolev spaces and conclude for the closedness of the solution mapping defined on the set of parameters. The second one contains a result on the behavior of Nash equilibrium points.

We start our exposition from the simple remark: if f is an upper semi-continuous function then the solution set of the inequality $f(x) \geq 0$ is closed.

Let X be a Hausdorff topological space and let $f_n : X \to \mathbb{R}$, $n \in \mathbb{N}$, be given functions. Let us consider, for $n \in \mathbb{N}^*$, the inequations

$$(1)_n \qquad f_n(x) \geq 0.$$

Denote by $S_1(1/n, f_n)$ the solutions set of $(1)_n$ and by $S_1(0, f_0)$ the solutions set of

$$(1) \qquad f_0(x) \geq 0.$$

Suppose that solutions sets are not empty for all $n \in \mathbb{N}$.

Let $M = \{f : X \to \mathbb{R} \,|\, \sup_{x \in X} |f(x)| < +\infty\}$ and let ρ be the metric on M given by

$$\rho(f, g) = \sup_{x \in X} |f(x) - g(x)|,\ f, g \in M.$$

*"Petru Maior" University, ROU - Târgu-Mureş, Romania, e-mail: bgdmarcel@netscape.net

†Department of Mathematics, Technical University of Cluj-Napoca, ROU - Cluj-Napoca, Romania, e-mail: Daniela.Inoan@math.utcluj.ro

BDD-A23788 © 2007 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 09:11:18 UTC)

We shall denote by $f_n \xrightarrow{\rho} f_0$ if $(f_n)_{n\in\mathbb{N}^*}$ converges (uniformly) to f_0 with respect to ρ.

Definition 1. *Let M' be a subset of M. We say that $(1)_n$ are stable if the solution mapping $S_1 : \mathbb{N}\times M' \to 2^X$ defined above is closed, i.e. if for each sequence $(x_n)_{n\in\mathbb{N}^*}$, $x_n \in S_1(1/n, f_n)$ with $x_n \to x$, $f_n \xrightarrow{\rho} f_0$ as $n \to +\infty$ one has $x \in S_1(0, f_0)$.*

If $M' = uC(X) = \{f \in M \mid f$ is upper semi-continuous on $X\}$ then $(1)_n$ are stable.

Concerning this property there are several results. Lignola and Morgan established convergence of the solutions of generalized parametric variational inequalities (see [6]). Related to Muu's article [7], more recently, Khanh and Luu [4], Li et al. [5] established *closedness* of the solutions function defined on the set of parameters for the same class of problems, namely parametric vector quasivariational inequalities. They have imposed conditions of upper semi-continuity being motivated by restrictions occurring in economical field. However, their hypotheses are too strong in some applications like variational inequalities governed by differential operators (see [2]).

2 Parametric equilibrium problems

Let X be a normed space, D a nonempty subset of X and let $f_n : X \times X \to \mathbb{R}$ be given functions. For $n \in \mathbb{N}^*$ we consider the following parametric equilibrium problem:

$$(2)_n \text{ Find an element } x_n \in D \text{ such that } f_n(x_n, y) \geq 0, \forall\ y \in D.$$

Denote by $S_2(1/n, f_n)$ the solutions set of $(2)_n$ and by $S_2(0, f_0)$ the solutions set of the "limit" equilibrium problem:

$$(2) \text{ Find an element } x_0 \in D \text{ such that } f_0(x_0, y) \geq 0, \forall\ y \in D.$$

Suppose that solutions sets are not empty for all $n \in \mathbb{N}$.

To compensate for the lack of upper semi-continuity, Brézis ([3]) introduced the notion of topological pseudomonotonicity (which is a kind of conditioned upper semi-continuity) only in the context of variational inequalities (see [8]).

Definition 2. *We say that a function $g : X \times X \to \mathbb{R}$ is* topologically pseudomonotone *(in the sense of Brézis) if for each sequence $(x_n)_{n\in\mathbb{N}}$ with $x_n \rightharpoonup x$ in X, $\liminf_n g(x_n, x) \geq 0$ imply*

$$\limsup_n g(x_n, y) \leq g(x, y), \text{ for all } y \in X.$$

Let $N = \{f : X \times X \to \mathbb{R} \mid \sup_{(x,y)\in X\times X} |f(x, y)| < +\infty\}$ and let d be the metric on N given by

$$d(f, g) = \sup_{(x,y)\in X\times X} |f(x, y) - g(x, y)|,\ f, g \in N.$$

BDD-A23788 © 2007 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 09:11:18 UTC)

We shall denote by $f_n \xrightarrow{d} f_0$ if $(f_n)_{n\in\mathbb{N}^*}$ converges (uniformly) to f_0 with respect to d.

Denote $pC(X,X) = \{f \in N \mid f \text{ is topologically pseudomonotone }\}$.

Proposition 1. *Let $(f_n)_{n\in\mathbb{N}^*}$ be a sequence such that $f_n \in pC(X,X)$. If $f_n \xrightarrow{d} f_0$ then $f_0 \in pC(X,X)$.*

Proof. Let $\varepsilon > 0$. Let $(x_n)_{n\in\mathbb{N}}$ with $x_n \rightharpoonup x$ in X, and $\liminf_n f_0(x_n,x) \geq 0$. For n large we gain $\liminf_n f_n(x_n,x) \geq 0$. Since f_n are in $pC(X,X)$ one has

$$\limsup_n f_n(x_n,y) \leq f_n(x,y), \text{ for all } y \in X.$$

Thus

$$\limsup_n f_0(x_n,y) \leq f_n(x,y) < f_0(x,y) + \varepsilon, \text{ for all } y \in X.$$

Letting $\varepsilon \to 0$ the conclusion follows. □

Definition 3. *Let N' be a subset of N. We say that $(2)_n$ are stable if the solution mapping $S_2 : \mathbb{N}\times N' \to 2^X$ defined above is closed, i.e. if for each sequence $(x_n)_{n\in\mathbb{N}^*}$, $x_n \in S_2(1/n, f_n)$ with $x_n \rightharpoonup x_0$, $f_n \xrightarrow{d} f_0$ as $n \to +\infty$ one has $x_0 \in S_2(0,f_0)$.*

Proposition 2. *If $N' = pC(X,X)$ then $(2)_n$ are stable.*

Proof. The conclusion follows by the following inequalities

$$\begin{aligned} 0 \leq \liminf_n f_n(x_n,y) &\leq \limsup_n f_0(x_n,y) + \liminf_n \big(f_n(x_n,y) - f_0(x_n,y)\big) \\ &= \limsup_n f_0(x_n,y) \leq f_0(x,y),\ \forall\, y \in D. \end{aligned}$$

□

3 Parametric Domains

To this point the feasible set has been "constant". Corresponding to $(1)_n$ for $n \in \mathbb{N}^*$ we shall consider the parametric inequations:

$$(1)'_n \qquad \text{find } x \in D_n \text{ such that } f_n(x) \geq 0,$$

where $D_n \subseteq X$ are nonempty and closed.

Denote by $S'_1(1/n, f_n)$ the solutions set of $(1)'_n$ and by $S'_1(0,f_0)$ the solutions set of

$$(1)' \qquad \text{find } x \in D_0 \text{ such that } f_0(x) \geq 0, x \in D_0.$$

Suppose that solutions sets are not empty for all $n \in \mathbb{N}$. Clearly $S'_1(1/n,f_n) = f_n^{-1}([0,+\infty)) \cap D_n$ and $S'_1(0,f_0) = f_0^{-1}([0,+\infty)) \cap D_0$. If $\emptyset \neq \cap_{n\in\mathbb{N}^*} D_n \subseteq D_0$ we can easily conclude that $(1)'_n$ are stable in the sense of Definition 1.

Corresponding to $(2)_n$ we shall consider the parametric equilibrium problems:

BDD-A23788 © 2007 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 09:11:18 UTC)

$(2)'_n$ Find an element $x_n \in D_n$ such that $f_n(x_n, y) \geq 0, \forall\ y \in D_n$.

Denote by $S'_2(1/n, f_n)$ the solutions set of $(2)'_n$ and by $S'_2(0, f_0)$ the solutions set of the "limit" equilibrium problem:

$(2)'$ Find an element $x_0 \in D_0$ such that $f_0(x_0, y) \geq 0, \forall\ y \in D_0$.

Suppose that solutions sets are not empty for all $n \in \mathbb{N}$.

Definition 4. *We say that $(2)'_n$ are stable if the solution map $S'_2 : \mathbb{N} \times pC(X, X) \to 2^X$ is closed, i.e. if $x_n \in S'_2(1/n, f_n)$, $x_n \rightharpoonup x_0$, and $f_n \xrightarrow{d} f_0$ in $pC(X, X)$ as $n \to +\infty$, then $x_0 \in S'_2(0, f_0)$.*

We have the following statement.

Theorem 1. *Let X be a normed space. Suppose that D_n are weakly closed for all $n \in \mathbb{N}^*$ and $\emptyset \neq \cap_{n\in\mathbb{N}^*} D_n = D_0$.*

Then $(2)'_n$ are stable in the sense of Definition 4.

Proof. Let $x_n \in D_n$ be such that $(2)'_n$ is valid. Let $x_n \rightharpoonup x_0$ in X thus we have $x_0 \in D_n$, so $\liminf_n f_n(x_n, x_0) \geq 0$. Since $f_n \xrightarrow{d} f_0$ it follows

$$\liminf_n f_0(x_n, x_0) \geq 0$$

therefore, by topological pseudomonotonicity of f_0 one has

$$\limsup_n f_0(x_n, y) \leq f_0(x_0, y), \text{ for all } y \in D_0.$$

Finally,

$$\begin{aligned} 0 \leq \liminf_n f_n(x_n, y) &\leq \limsup_n f_0(x_n, y) + \liminf_n \big(f_n(x_n, y) - f_0(x_n, y)\big) \leq \\ &\leq \limsup_n f_0(x_n, y) \leq f_0(x_0, y),\ \forall\, y \in D_0, \end{aligned}$$

which completes the proof. □

One question rises. What is the condition for the parametric domains in order to have the stability of $(2)'_n$? We are not naive to consider condition $\emptyset \neq \cap_{n\in\mathbb{N}^*} D_n = D_0$ a proper one, especially in applications. In [2] we used Mosco's convergence.

4 Application to hemivariational inequalities

Let us denote by $pC_0(X, X) = \{f \in pC(X, X) : f(x, x) = 0, x \in X\}$. First, we establish that $pC_0(X, X)$ is stable with respect to addition. The proof is similar to Proposition 2.4 in [8].

Proposition 3. *If $f, g \in pC_0(X, X)$ then $f + g \in pC_0(X, X)$.*

BDD-A23788 © 2007 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 09:11:18 UTC)

Proof. We claim that $\liminf_n f(x_n, x) \geq 0$ and $\liminf_n g(x_n, x) \geq 0$. Otherwise $\liminf_n g(x_n, x) = -\varepsilon < 0$ and, by passing to a subsequence $\lim_n g(x_n, x) = -\varepsilon$. Therefore

$$\begin{aligned}\liminf_n f(x_n, x) &= \liminf_n \big((f+g)(x_n, x) - g(x_n, x)\big) \\ &= \liminf_n (f+g)(x_n, x) - \lim_n g(x_n, x) \geq 0 + \varepsilon.\end{aligned}$$

Since f is topologically pseudomonotone one has

$$\limsup_n f(x_n, y) \leq f(x, y), \text{ for all } y \in X.$$

Now, put $x = y$ to reach the contradiction as follows

$$\varepsilon \leq \liminf_n f(x_n, x) \leq \limsup_n f(x_n, x) \leq f(x, x) = 0.$$

The conclusion is obtained from the super-additivity of lim inf and sub-additivity of lim sup . □

Let Ω be a bounded open set of $\mathbb{R}^N$ with Lipschitz boundary. Let us consider a function $l : \Omega \times \mathbb{R} \times \mathbb{R} \to \mathbb{R}$, such that $l(x; \cdot, \cdot)$ is upper semi-continuous for a.e. $x \in \Omega$ and $l(\cdot; y, z)$ is measurable for all $y, z \in \mathbb{R}$. We say that l satisfies *the growth condition* if there exist $h_1, h_2 \in \mathbb{R}_+$ such that

$$|l(x; y, z)| \leq (h_1 + h_2|y|)|z|, \text{ a.e. } x \in \Omega, \forall\, y, z \in \mathbb{R}.$$

Lemma 1. ([2]) *If l satisfies the conditions above, then the function defined on $H^1(\Omega) \times H^1(\Omega)$ by*

$$(u, w) \rightarrowtail \int_\Omega l\big(x; u(x), w(x)\big)\, dx$$

is weakly upper semi-continuous.

For $n \in \mathbb{N}^*$ we consider the following variational inequality problem:
$(VI)_n$ Find an element $u_n \in K$ such that

$$\begin{aligned}&\int_\Omega A_n\big(x, u_n(x), \nabla u_n(x)\big) \cdot \big(\nabla v(x) - \nabla u_n(x)\big)\, dx \; + \\ &\quad + \int_\Omega a^n\big(x, u_n(x), \nabla u_n(x)\big)\big(v(x) - u_n(x)\big)\, dx \; + \\ &\quad + \int_\Omega l_n\big(x; u_n(x), v(x) - u_n(x)\big)\, dx \geq 0, \forall\, v \in K,\end{aligned}$$

with $A_n = (a_1^n, \ldots, a_N^n)$, a^n, and l_n given functions, and K a closed, convex, nonempty subset of the Sobolev space $H^1(\Omega)$.

For a fixed $n \in \mathbb{N}$, we consider the operator

$$\mathcal{A}_n : H^1(\Omega) \to (H^1(\Omega))^*, \qquad \mathcal{A}_n(u) = \mathcal{B}_n(u, u),$$

BDD-A23788 © 2007 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 09:11:18 UTC)

where

$$\mathcal{B}_n(u,v)(w) = \int_\Omega \Big\{ \sum_{i=1}^{N} a_i^n(x,u(x),\nabla u(x))\cdot\partial_i w(x)\Big\}dx + \int_\Omega a^n(x,u(x),\nabla u(x))w(x)dx.$$

The operators $\mathcal{A}_n$ are topologically pseudomonotone if satisfies some reasonable conditions (see [8], page 76; [2]).

$(VI)_n$ can be written as

$$\text{find } u_n \in K \text{ such that } f_n(u_n,v) + L_n(u_n,v) \geq 0, \forall\ v \in K,$$

where $f_n(u,v) = \langle \mathcal{A}_n(u), v-u\rangle$ and $L_n(u,v) = \displaystyle\int_\Omega l_n\big(x; u(x), v(x)-u(x)\big)\,dx$.

Finally, we can apply Proposition 2 and Proposition 3 to obtain a result on the stability of $(VI)_n$.

References

[1] Anh, L. Q., and Khanh, P. Q.: *Uniqueness and Hölder continuity of the solution to multivalued equilibrium problems in metric spaces*, Journal of Global Optimization, original paper, DOI 10.1007/s10898-006-9062-8 (2006).

[2] Bogdan, M., and Kolumbán, J.: *On Nonlinear Parametric Variational Inequalities*, Nonlinear Analysis, Vol. 67, No. 7, pp. 2272 - 2282, 2007.

[3] Brézis, H.: *Équations et inéquations non linéaires en dualité*, Annales de l'Institut Fourier (Grenoble) 18(1)(1965), pp. 115 - 175.

[4] Khanh, P. Q., and Luu, L. M.: *Upper Semicontinuity of the Solution Set to Parametric Vector Quasivariational Inequalities*, Journal of Global Optimization, **32**(2005), 569 - 580.

[5] Li, S. J., Chen, G. Y., and Teo, K. L.: *On the Stability of Generalized Vector Quasivariational Inequality Problems*, Journal of Optimization Theory and Applications, vol. 107, no. 1, pp. 35 - 50, 2000.

[6] Lignola, M. B., and Morgan, J.: *Generalized Variational Inequalities with Pseudomonotone Operators Under Perturbations*, Journal of Optimization Theory and Applications, Vol. 101, No. 1, pp. 213 - 220, 1999.

[7] Muu, L. D.: *Stability property of a class of variational inequalities*, Mathematische Operationforschung und Statistik, Serie Optimization 15, pp. 347 - 351, 1984.

[8] Showalter, R. E.: Monotone Operators in Banach Space and Nonlinear Partial Differential Equations, American Mathematical Society, vol. 49, 1997.

[9] Yu, J., Yang, H., and Yu, C.: *Well-posed Ky Fan's point, quasi-variational inequality and Nash equlibrium problems*, Nonlinear Analysis, Vol. 66, pp. 777 - 790, 2007.

BDD-A23788 © 2007 Editura Universităţii „Petru Maior"
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 09:11:18 UTC)