# REPRESENTATIONS OF POSITIVE INTEGERS AS SUM OF SQUARES AND TRIANGULAR NUMBERS

R.Y. Denis [*], [†] S.N. Singh [‡] and S.P. Singh [§¶]

Proceedings of the European Integration between Tradition and Modernity
"Petru Maior" University of Târgu-Mureş
Oktober, 22-23, 2009

**Abstract**

In this paper, using some known summations, we have established the generating functions for the number of representations of a positive integer as the sum of squares of integers and also as the sum of triangular numbers.

*Key words and phrases:* Sum of squares, sum of triangular numbers, general theta function.

*2000 AMS Subject classification:* Primary: 11E25, 33E05; Secondary: 05A15, 33D15

## 1 Introduction, notations and definitions

One of the most interesting problems in number theory is the representation of positive integers as sum of squares of integers.Fermat proved that all primes of the form $4n + 1$can be,uniquely, expressed as the sum of two

[*]Department of Mathematics, University of Gorakhpur, Gorakhpur-273009, India
[†]Corresponding author
[‡]Department of Mathematics, University of Gorakhpur, Gorakhpur-273009, India
[§]Department of Mathematics, T.D.P.G. College, Jaunpur-222002, India
[¶]E-mail: ddry@rediffmail.com



BDD-A23666 © 2009 Editura Universităţii „Petru Maior"
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 22:58:03 UTC)

squares. Lagrange showed that all positive integers can be represented as sum of four squares and that this number is minimal.
Let $\gamma_k(n)$ stand for the number of ways the positive integer $n$ can be represented as a sum of $k$ squares with representations arising from different signs and from different orders being regarded as distinct.
Since,

$$1 = (\pm 1)^2 + 0^2 = 0^2 + (\pm 1)^2,$$

we have $\gamma_2(1) = 4$. Also,

$$1 = (\pm 1)^2 + 0^2 + 0^2 = 0^2 + (\pm 1)^2 + 0^2 = 0^2 + 0^2 + (\pm 1)^2$$

so, we have $\gamma_3(1) = 6$.
Ramanujan's general theta function $f(a, b)$ is defined by

$$f(a,b) = \sum_{n=-\infty}^{\infty} a^{n(n+1)/2} b^{n(n-1)/2}, \ |ab| < 1.$$

The most important special cases of $f(a, b)$ are given by,

$$\phi(q) = f(q,q) = \sum_{n=-\infty}^{\infty} q^{n^2} = \frac{[-q;-q]_\infty}{[q;-q]_\infty} = \frac{[q^2,-q;q^2]_\infty}{[-q,q;q^2]_\infty},$$

$$\psi(q) = f(q,q^3) = \sum_{n=0}^{\infty} q^{n(n+1)/2} = \frac{[q^2;q^2]_\infty}{[q;q^2]_\infty}, \tag{1}$$

where $[\alpha;q]_\infty = \Pi_{r=0}^{\infty}(1-\alpha q^r)$.
The generating function for $\gamma_k(n)$ is given by

$$\sum_{n=0}^{\infty} \gamma_k(n) q^n = \phi^k(q) = \Big\{ \sum_{n=-\infty}^{\infty} q^{n^2} \Big\}^k.$$

The well known two and four square theorems due to Jacobi are:

$$\gamma_2(n) = 4[d_1(n) - d_3(n)]$$

and

$$\gamma_4(n) = 8 \sum_{d/n;\& \ 4 \ does \ not \ divide \ n} d,$$



BDD-A23666 © 2009 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 22:58:03 UTC)

where $d_i(n)$ denotes the number of positive divisors of $n$ that are congruent to $i$ (modulo 4).
A triangular number is a number of the form $n(n+1)/2$ for non negative integer $n$. Let $t_k(n)$ denote the number of representations of $n$ as the sum of $k$ triangular numbers, then generating function for $t_k(n)$ is given by

$$\sum_{n=0}^{\infty} t_k(n)q^n = \psi^k(q) = \Big\{ \sum_{j=0}^{\infty} q^{j(j+1)/2} \Big\}^k,$$

where $\psi(q)$ is given by (1).
The first few triangular numbers are $0, 1, 3, 6, 10, 15, ...$ Thus, $t_2(6) = 3$.
Gauss proved that every natural number is the sum of 3 or fewer triangular numbers. Ewell [4] used Jacobi's triple product identity to show that,

$$t_2(n) = d_1(4n+1) - d_3(4n+1).$$

Adiga [1] used ${}_1\psi_1$ summation formula due to Ramanujan to show that

$$t_4(n) = \sum_{d/(2n+1)} d.$$

Coper and Lam [3], using the summation formula for ${}_1\psi_1$ derived formulas for $t_2(n), t_4(n), t_6(n)$, and $t_8(n)$. We produce here $t_6(n)$ and $t_8(n)$,

$$t_6(n) = \frac{1}{8} \sum_{d/4n+3} (-1)^{\{(d+1)/2\}} d^2$$

and

$$t_8(n) = \sum_{d/(n+1), d-odd} \Big\{ \frac{(n+1)}{d} \Big\}^3.$$

We define the basic bilateral hypergeometric series as

$${}_r\psi_r \left[ \begin{matrix} a_1, a_2, ..., a_r; \\ b_1, b_2, ..., b_r \end{matrix} q; \; z \right] = \sum_{n=-\infty}^{\infty} \frac{[a_1, a_2, ..., a_r]_n \, z^n}{[b_1, b_2, ..., b_r]_n},$$

valid for $\max\{|a_1|, |a_2|, ..., |a_r|, |b_1|, |b_2|, ..., |b_r|, |z|\} < 1$.



BDD-A23666 © 2009 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 22:58:03 UTC)

## 2 Main Results

In this section we shall establish certain identities from some known summations and then use these to derive the values of $\gamma_k(n)$ and $t_k(n)$.

(a) Ramanujan's remarkable summation formula for ${}_1\psi_1$ is

$$ {}_1\psi_1\left[\begin{array}{c} a;\ q;\ z \\ b \end{array}\right] = \sum_{n=0}^{\infty} \frac{[a;q]_n z^n}{[b;q]_n} = \frac{[az, q/az, q, b/a; q]_\infty}{[z, b/az, b, q/a; q]_\infty}. \tag{2} $$

Now, replacing $q$ by $q^k$ and then setting $a = q^j, b = q^{k+j}$ and $z = q^i$ in (2), we get

$$ \sum_{n=-\infty}^{\infty} \frac{q^{in}}{1-q^{kn+j}} = \frac{[q^k;q^k]_\infty^2 [q^{i+j}, q^{k-i-j}; q^k]_\infty}{[q^i, q^j, q^{k-i}, q^{k-j}; q^k]_\infty} \tag{3} $$

provided $i, j \not\equiv 0\ mod(k)$.

(i) Now, taking $k = 4$, $i = j = 1$ in (3), we find

$$ \begin{aligned} \sum_{n=-\infty}^{\infty} \frac{q^n}{1-q^{4n+1}} &= \sum_{n=0}^{\infty} \frac{q^n}{1-q^{4n+1}} - \sum_{n=0}^{\infty} \frac{q^{3n+2}}{1-q^{4n+3}} = \frac{[q^2;q^2]_\infty^2}{[q;q^2]_\infty^2} = \psi^2(q) \\ &= \sum_{n=0}^{\infty} t_2(n) q^{2n} \end{aligned} \tag{4} $$

One can notes that (4) implies the following well known identity

$$ t_2(n) = d_1(4n+1) - d_3(4n+1). $$

(ii) Again, taking $k = 4$, $i = 2$ and $j = 1$ in (3), we get

$$ \sum_{n=-\infty}^{\infty} \frac{q^{2n}}{1-q^{4n+1}} = \frac{[q^4;q^4]_\infty^2}{[q^2;q^4]_\infty^2} = \psi^2(q^2). \tag{5} $$

Thus we have

$$ \sum_{n=0}^{\infty} \frac{q^{2n}}{1-q^{4n+1}} - \sum_{n=0}^{\infty} \frac{q^{2n+1}}{1-q^{4n+3}} = \psi^2(q^2) = \sum_{n=0}^{\infty} t_2(n) q^{2n}. \tag{6} $$



BDD-A23666 © 2009 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 22:58:03 UTC)

Now, equating the coefficients of $q^{2n}$ on both sides of (6), we get

$$t_2(n) = d_1(4n+1) - d_3(4n+1).$$

Ramanujan's ${}_1\psi_1$ summation can, also, be written as

$$\sum_{n=0}^{\infty} \frac{[q/\beta;q]_n(-\beta q^{-1/2}/z)^n}{[\alpha q;q]_n} + \sum_{n=1}^{\infty} \frac{[1/\alpha;q]_n(-\alpha z q^{1/2})^n}{[\beta;q]_n} = \frac{[-zq^{1/2}, -q^{1/2}, q, \alpha\beta;q]_\infty}{[-\alpha z q^{1/2}, -\beta/zq^{1/2}, \alpha q, \beta;q]_\infty}. \quad (7)$$

Differentiating both sides of (7) with respect to $z$ and then setting $z = q^{-1/2}$ Bhargava and Somashekara [2] established the following identity,

$$\sum_{n=0}^{\infty} \frac{(n+1)[q/\alpha;q]_n\alpha^n}{[\beta;q]_{n+1}} + \sum_{n=0}^{\infty} \frac{n[q/\beta;q]_n\beta^n}{[\alpha;q]_{n+1}} = \frac{[q;q]_\infty^3[\alpha\beta;q]_\infty}{[\alpha,\beta;q]_\infty^2}. \quad (8)$$

Replacing $q$ by $q^k$ and then taking $\alpha = q^i$ and $\beta = q^j$ in (8), we get

$$\sum_{n=0}^{\infty} \frac{(n+1)[q^{k-i};q^k]_n q^{in}}{[q^j;q^k]_{n+1}} + \sum_{n=0}^{\infty} \frac{n[q^{k-j};q^k]_n q^{jn}}{[q^i;q^k]_{n+1}} = \frac{[q^k;q^k]_\infty^3[q^{i+j};q^k]_\infty}{[q^i,q^j;q^k]^2\infty}; \quad (i,j,i+j<k). \quad (9)$$

Now, taking $k=2$, $i=j=1$ in (9), we get

$$\sum_{n=0}^{\infty} \frac{(2n+1)q^n}{1-q^{2n+1}} = \frac{[q^2;q^2]_\infty^4}{[q;q^2]_\infty^4} = \psi^4(q) = \sum_{n=0}^{\infty} t_{4(n)}q^n$$

which has the arithmetic consequence

$$t_4(n) = \sum_{d/(2n+1)} d,$$

which is a known result due to Legendre.

Ramanujan [6] in Chapter 17, page 139, of the second notebook has mentioned the following identity,



BDD-A23666 © 2009 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 22:58:03 UTC)

$$\sum_{n=1}^{\infty} \frac{n^3 q^{n-1}}{1-q^{2n}} = \psi^8(q),$$

which can be written as

$$\sum_{n=0}^{\infty} \frac{(n+1)^3 q^n}{1-q^{2n+2}} = \psi^8(q) = \sum_{n=0}^{\infty} t_8(n) q^n$$

which the following arithmetic interpretation,

$$t_{8(n)} = \sum_{d/n+1, d-odd} \left\{\frac{n+1}{d}\right\}^3.$$

(b) Bailey's sum of a well-possed ${}_3\psi_3$ is

$$\sum_{n=-\infty}^{\infty} \frac{[b,c,d;q]_n (q/bcd)^n}{[q/b,q/c,q/d;q]_n} = \frac{[q,q/bc,q/bd,q/cd;q]_\infty}{[q/b,q/c,q/d,q/bcd;q]_\infty}$$

which can be re-written as,

$$1 + \sum_{n=1}^{\infty} \frac{[b,c,d;q]_n (1+q^n)(q/bcd)^n}{[q/b,q/c,q/d;q]_n} = \frac{[q,q/bc,q/bd,q/cd;q]_\infty}{[q/b,q/c,q/d,q/bcd;q]_\infty}. \quad (10)$$

The above relation can also be deduced from the well known basic bilateral analogue of Dixon's summation, (cf. Gasper-Rahman [5]; App.II(II.32)]). Now, taking $b = c = d = -1$ in (10), we get

$$1 + 8\sum_{n=1}^{\infty} \frac{(-1)^n q^n}{(1+q^n)^2} = \frac{[q;q]_\infty^4}{[-q;q]_\infty^4} = \phi^4(-q). \quad (11)$$

Now, replacing $q$ by $-q$ in (11), we have

$$1 + 8\sum_{n=1}^{\infty} \frac{q^n}{\{1+(-q)^n\}^2} = \phi^4(q). \quad (12)$$



BDD-A23666 © 2009 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 22:58:03 UTC)

Now, (12) can be interpreted as

$$\begin{aligned}
\phi^4(q) &= \sum_{n=0}^{\infty}\gamma_4(n)q^n = 1+8\sum_{n=1}^{\infty}\frac{q^n}{\{1+(-q)^n\}^2} = 1+8\sum_{n=1}^{\infty}\frac{nq^n}{1+(-q)^n}\\
&= 1+8\Big\{\sum_{n\geq 1,n-odd}\frac{nq^n}{1-q^n}+\sum_{n\geq 2,n-even}\frac{nq^n}{1+q^n}\Big\}\\
&= 1+8\Big\{\sum_{n\geq 1}\frac{nq^n}{1-q^n}+\sum_{n\geq 2,n-even}\Big(\frac{nq^n}{1+q^n}-\frac{nq^n}{1-q^n}\Big)\Big\}\\
&= 1+8\Big\{\sum_{n=1}^{\infty}\frac{nq^n}{1-q^n}-\sum_{n\geq 2,n-even}\frac{2nq^{2n}}{1-q^{2n}}\Big\}\\
&= 1+8\Big\{\sum_{n=1}^{\infty}\frac{nq^n}{1-q^n}-\sum_{n=1}^{\infty}\frac{4nq^{4n}}{1-q^{4n}}\Big\}
\end{aligned}$$

which implies that

$$1+\sum_{n=1}^{\infty}\gamma_{4(n)}q^n = 1+8\sum_{4\ does\ not\ divide\ n}\frac{nq^n}{1-q^n}. \tag{13}$$

Now, equating the coefficients of $q^n$ on both sides of (13), we get,

$$\gamma_{4(n)} = 8\sum_{d\ divides\ n\ but\ 4\ does\ not\ divide\ n} d.$$

Again, letting $d\to\infty$ in (10), we get

$$1+\sum_{n=1}^{\infty}\frac{(-1)^n q^{n(n-1)/2}(1+q^n)[b,c;q]_n(q/bc)^n}{[q/b,q/c]_n} = \frac{[q,q/bc;q]_\infty}{[q/b,q/c;q]_\infty}. \tag{14}$$

Now, setting $b=c=-1$ in (14) and replacing $q$ by $-q$ in it, we get

$$1+4\sum_{n=1}^{\infty}\frac{(-1)^{n(n-1)/2}q^{n(n+1)/2}}{1+(-q)^n} = \frac{[-q;-q]_\infty^2}{[q;-q]_\infty^2} = \phi^2(q). \tag{15}$$



BDD-A23666 © 2009 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 22:58:03 UTC)

Thus, we have

$$\begin{aligned}\phi^2(q) &= \sum_{n=0}^{\infty}\gamma_2(n)q^n = 1+4\sum_{n=1}^{\infty}\Big\{\frac{q^{4n-3}}{1-q^{4n-3}}-\frac{q^{4n-1}}{1-q^{4n-1}}\Big\}\\ &= \sum_{n=0}^{\infty}\gamma_2(n)q^n = 1+4\sum_{n=0}^{\infty}\Big\{\frac{q^{4n+1}}{1-q^{4n+1}}-\frac{q^{4n+3}}{1-q^{4n+3}}\Big\}.\end{aligned}$$

Now, equating the coefficients of $q^n$ on both sides of the above relation, we get Jacobi's two square theorem,

$$\gamma_2(n) = 4\{d_1(n) - d_3(n)\},$$

where $d_i(n)$ is the number of positive divisors of $n$ that are congruent to $i$ (*modulo* 4).

(c) The basic bilateral, (cf.Gasper-Rahman [5]; App.II(II.32)), analogue of Dixon's summation is

$$\sum_{n-\infty}^{\infty}\frac{(1+q^n a^{1/2})[b,c,d;q]_n(qa^{3/2})^n}{(1+a^{1/2})[aq/b,aq/c,aq/d;q]_n(bcd)^n}$$
$$= \frac{[aq,aq/bc,aq/bd,aq/cd,qa^{1/2}/b,qa^{1/2}/c,qa^{1/2}/d,q,q/a;q]_\infty}{[aq/b,aq/c,aq/d,q/b,q/c,q/d,qa^{1/2},q/a^{1/2},qa^{3/2}/bcd;q]_\infty}.$$

Letting $a \to 1$ in (5), we get

$$1+\sum_{n=1}^{\infty}\frac{[b.c.d;q]_n(1+q^n)(q/bcd)^n}{[q/b,q/c,q/d;q]_n} = \frac{[q,q/bc,q/bd,q/cd;q]_\infty}{[q/b,q/c,q/d,q/bcd;q]_\infty}$$

which is (10). Now proceeding on the above lines we can again find the identities (12) and (15) from which $\gamma_4(n)$ and $\gamma_2(n)$ can be formulated. In entry 25 of Chapter 16 of second Notebook, Ramanujan has mentioned the following beautiful identities:

$$\phi(q) + \phi(-q) = 2\phi(q^4)$$

$$\phi(q) - \phi(-q) = 4q\psi(q^8)$$



BDD-A23666 © 2009 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 22:58:03 UTC)

$$\phi(q)\phi(-q) = \phi^2(-q^2) \tag{16}$$

$$\psi(q)\psi(-q) = \psi(q^2)\phi(-q^2)$$

$$\phi(q)\psi(q^2) = \psi^2(q) \tag{17}$$

$$\phi^2(q) - \phi^2(-q) = 8q\psi^2(q^4) \tag{18}$$

$$\phi^2(q) + \phi^2(-q) = 2\phi^2(q^2) \tag{19}$$

$$\phi^4(q) - \phi^4(-q) = 16q\phi^4(q^2). \tag{20}$$

Now, from (20) we have

$$\sum_{k=0}^{\infty} \gamma_4(k)q^k - \sum_{k=0}^{\infty} \gamma_4(k)(-q)^k = 16\sum_{k=0}^{\infty} \gamma_4(k)q^{2k+1}. \tag{21}$$

Now, equating the coefficients of $q^{2k+1}$ on both sides of (21), we get

$$\gamma_4(2k+1) = 8\gamma_4(k).$$

Further, from (19) we have

$$\sum_{k=0}^{\infty} \gamma_2(k)q^k + \sum_{k=0}^{\infty} \gamma_2(-q)^k = 2\sum_{k=0}^{\infty} \gamma_2(k)q^{2k}. \tag{22}$$

Now, equating the coefficients of $q^{2k}$ on both sides of (22), we get $\gamma_2(2k) = 2\gamma_2(k)$. Again, (18) leads to

$$\sum_{k=0}^{\infty} \gamma_2(k)q^k - \sum_{k=0}^{\infty} \gamma_2(k)(-q)^k = 8\sum_{k=0}^{\infty} t_2(k)q^{4k+1}$$



BDD-A23666 © 2009 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 22:58:03 UTC)

which, on equating the coefficients of $q^{4k+1}$ on both sides, leads to

$$\gamma_2(4k+1) = 4t_2(k).$$

Squaring both sides of (17) we get

$$\phi^2(q)\psi^2(q^2) = \psi^4(q)$$

which leads to

$$\sum_{m=0}^{\infty} \gamma_2(m)q^m \sum_{n=0}^{\infty} t_2(n)q^{2n} = \sum_{k=0}^{\infty} t_4(k)q^k. \tag{23}$$

Now, equating the coefficients of $q^{2k}$ on both sides of (23), we get

$$\sum_{n=0}^{k} \gamma_2(2k-2n)t_2(n) = t_4(2k).$$

Again, squaring both sides of (16), we get

$$\phi^2(q)\phi^2(-q) = \phi^4(-q^2),$$

which can be put in the form

$$\sum_{m=o}^{\infty} \gamma_2(m)q^m \sum_{n=0}^{\infty} \gamma_2(n)(-q)^n = \sum_{k=0}^{\infty} \gamma_4(k)(-1)^k q^{2k}.$$

Now, setting $m = 2k-n$ in the above and equating the coefficients of $q^{2k}$ on both sides,we get

$$\sum_{n=0}^{2k} (-1)^n \gamma_2(n)\gamma_2(2k-n) = (-1)^k \gamma_4(k)$$

which is the same as,

$$\sum_{n=0}^{4k} (-1)^n \gamma_2(n)\gamma_2(4k-n) = \gamma_4(2k).$$

Similarly, scores of other relations can also be established.



BDD-A23666 © 2009 Editura Universităţii „Petru Maior”
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 22:58:03 UTC)

**Acknowledgement**. This work has been done under Major Research Projects F. 31-83/2005 (S.R.) and F.No. 32-164/2006(SR) sanctioned to the first two authors, respectively by the University Grants Commission, Govt. of India, New Delhi, for which they wish to thank the authorities concerned. The first two authors are also thankful to the Department of Science and Technology, Govt. of India, New Delhi, for support under major research projects No. SR/S4/MS-461/07 dtd. 13.2.2008 entitled "A study of basic hypergeometric functions with special reference to Ramanujan mathematics" and No.SR/S4/MS:524 dtd. 10.2.2008 entitled "Glimpses of Ramanujan's mathematics in the field of $q$-series" sanctioned to them, respectively.

# References

[1] Adiga, C., *Sums of squares and sums of triangular numbers and their relationship*, The Math. Student, 73(1-4)(2004)87-94.

[2] Bhargava, S. and Somashekara, D.D., *Some Eta function identities deduced from Ramanujan's ${}_1\psi_1$ summation*, J. Math. Analysis and Applications, 176(2)(1993)554-560.

[3] Cooper, S. and Lam, H.Y., *Sums of two, four, six and eight squares and triangular numbers: An elementary approach*, Indian J. Math., 44(1)(2002)21-40.

[4] Ewell, J.A., *On sums of triangular numbers and sums of squares,* Amer. Math. Monthly, 99(1992)752-757.

[5] Gasper, G. and Rahman, M., Basic Hypergeometric Series, Cambridge University Press (1991).

[6] Ramanujan, S., Ramanujan's Notebook Vol.II, T.I.F.R., Bombay (1957).



BDD-A23666 © 2009 Editura Universităţii „Petru Maior"
Provided by Diacronia.ro for IP 216.73.217.142 (2026-11-14 22:58:03 UTC)