On the functional equation of generalized pseudomediality

ADRIAN PETRESCU "Petru-Maior" University of Târgu-Mureş, Romania apetrescu@upm.ro

Abstract

Based on $D_{i,j}$ and D_{i-j} conditions (see [1]) we give a direct and straightforward method to solve the functional equation of generalized pseudomediality on quasigroups.

2000 Mathematics Subject Classification: 39B52, 20N05, 20N15.

In this paper we solve the functional equation of generalized pseudomediality on quasigroups

$$\alpha_1(\alpha_2(x,y),\alpha_3(z,u)) = \alpha_4(\alpha_5(\alpha_6(x,z),y)u) \tag{1}$$

where x, y, z, u are taken from an arbitrary set A and α_i are quasigroup operations on A.

Sokhats'kyi [3] proved that every quadratic parastrophically uncancelable functional equation for four object variables is parastrophically equivalent to the functional equation of mediality (for this equation see [2]) or the functional equation of pseudomediality. He solved this equation using the functional equation of generalized associativity (for this equation see [2]).

The method we used is an example of the application of our results developed in [1].

Let $\alpha_1, \ldots, \alpha_6$ be a six quasigroup operations on A and forming a solution of the functional equation (1). We define $\alpha: A^4 \to A$ by

$$\alpha(x, y, z, u) = \alpha_1(\alpha_2(x, y), \alpha_3(z, u)) = \alpha_4(\alpha_5(\alpha_6(x, z), y), u)$$
 (2)

It is obvious that (A, α) is 4-quasigroup and conditions $D_{1,2}$, $D_{3,4}$ and D_{1-3} are fulfilled in (A, α) (see [1]).

Theorem 1. In a 4-loop (A, α) condition $D_{1,2}\&D_{3,4}\&D_{1-3}$ holds iff $\alpha(x, y, z, u) = yxzu$ where (A, \cdot) is a binary group.

Proof. Suppose that in 4-loop (A, α) condition $D_{12}\&D_{3,4}\&D_{1-3}$ holds and let e be a unit in this loop. We define

$$x \cdot y = \alpha(e, e, x, y) \tag{3}$$

Then (A, \cdot) is a binary loop with the unit e. From

$$\alpha(e, e, z, u) = \alpha(e, e, e, \alpha(e, e, z, u))$$

by condition $D_{3,4}$ we obtain

$$\alpha(x, y, z, u) = \alpha(x, y, e, zu) \tag{4}$$

Now

$$\alpha(x, y, e, zu) = \alpha(e, y, zu) \quad \text{(condition } D_{1-3})$$

$$= \alpha(e, y, e, x(zu)) \quad \text{(by (4))}$$

$$= \alpha(y, e, e, x(zu)) \quad \text{(condition } D_{1,2})$$

$$= \alpha(e, e, y, x(zu)) \quad \text{(condition } D_{1-3})$$

$$= y(x(zu)). \quad \text{(by (4))}$$

Thus

$$\alpha(x, y, z, u) = y(x(zu)) \tag{5}$$

Putting z = u = e in (5) we get

$$\alpha(x, y, e, e) = yx \tag{6}$$

From

$$\alpha(x, y, e, e) = \alpha(\alpha(x, y, e, e), e, e, e)$$

by condition $D_{1,2}$ we have

$$\alpha(x, y, z, u) = \alpha(yx, e, z, u) \tag{7}$$

Now

$$\alpha(yx, e, z, u) = \alpha(e, yx, z, u) \quad \text{(condition } D_{1,2})$$

$$= \alpha(z, yx, e, u) \quad \text{(condition } D_{1-3})$$

$$= \alpha((yx)z, e, e, u) \quad \text{(by (7))}$$

$$= \alpha((yx)z, e, u, e) \quad \text{(condition } D_{3,4})$$

$$= \alpha(e, (yx)z, u, e) \quad \text{(condition } D_{1,2})$$

$$= \alpha(u, (yx)z, e, e) \quad \text{(condition } D_{1-3})$$

$$= ((yx)z)u \quad \text{(by (7))}$$

Thus

$$\alpha(x, y, z, u) = ((yx)z)u \tag{8}$$

From (5) and (8) we have

$$y(x(zu)) = (yx)z)u (9)$$

Putting u = e in (9) we obtain

$$y(xz) = (yx)z$$

Therefore (A, \cdot) is a group and $\alpha(x, y, z, u) = yxzu$ The converse is obvious. **Theorem 2.** The set of all solutions of the functional equation of generalized pseudomediality over the set of quasigroup operations on an arbitrary set A is described by the realtions

$$\alpha_{1}(x,y) = F_{1}(x) + F_{2}(y) \qquad \alpha_{2}(x,y) = F_{1}^{-1}(F_{3}(y) + F_{4}(x))
\alpha_{3}(x,y) = F_{2}^{-1}(F_{5}(x) + F_{6}(y)), \quad \alpha_{4}(x,y) = F_{7}(x) + F_{6}(y)
\alpha_{5}(x,y) = F_{7}^{-1}(F_{3}(y) + F_{8}(x)), \quad \alpha_{6}(x,y) = F_{8}^{-1}(F_{4}(x) + F_{5}(y))$$
(10)

where (A, +) is an arbitrary group and F_1, \ldots, F_8 are arbitrary substitutions of the set A

Proof. Let $\alpha_1, \ldots, \alpha_6$ be six quasigroup operations on A and forming a solution of equation (1). We define $\alpha: A^4 \to A$ by (2). From the above results if follows that $\alpha(x, y, z, u) = T_2(y) + T_1(x) + T_3(z) + T_4(u)$, where (A, +) is a group with zero element $0 = T_1(a_1) = T_2(a_2) = T_3(a_3) = T_4(a_4)$, T_i being translations by $a = (a_1^4) \in A^4$ in (A, α) (for details see [1] and [2]). Putting $z = a_3$ and $u = a_4$ in (2) we get

$$\alpha_1(\alpha_2(x,y),\alpha_3(a_3,a_4)) = \alpha_4(\alpha_5(\alpha_6(x,a_3),y),a_4) = T_2(y) + T_1(x). \tag{11}$$

The mappings $f(x) = \alpha_1(x, \alpha_3(a_3, a_4)), f_1(x) = \alpha_6(x, a_3)$ and $f_2(x) = \alpha_4(x, a_4)$ are substitutions of the set A. From (11) we obtain

$$\alpha_2(x,y) = f^{-1}(T_2(y) + T_1(x))$$

and

$$\alpha_5(x,y) = f_2^{-1}(T_2(y) + T_1(f_1^{-1}(x))).$$

For $x = a_1$ and $y = a_2$ in (2) we have

$$\alpha_1(\alpha_2(a_1, a_2), \alpha_3(z, u)) = \alpha_4(\alpha_5(\alpha_6(a_1, z), a_2), u) = T_3(z) + T_4(u)$$

and thus

$$\alpha_3(z,u) = g^{-1}(T_3(z) + T_4(u))$$

where

$$g(x) = \alpha_1(\alpha_2(a_1, a_2), x)$$

and

$$\alpha_4(z, u) = T_3(g_1^{-1}(z)) + T_4(u)$$

where

$$g_1(x) = \alpha_5(\alpha_6(a_1, x), a_2).$$

Finally, if we put $y = a_2$ and $u = a_4$ in (2) then we have

$$\alpha_1(\alpha_2(x, a_2), \alpha_3(z, a_4)) = \alpha_4(\alpha_5(\alpha_6(x, z), a_2), a_4) = T_1(x) + T_3(z)$$

and thus

$$\alpha_1(x,z) = T_1(h_1^{-1}(x)) + T_3h_2^{-1}(z)$$

for $h_1(x) = \alpha_2(x, a_2)$ and $h_2(x) = \alpha_3(x, a_4)$,

$$\alpha_6(x,z) = h^{-1}(T_1(x) + T_3(z))$$

where $h(x) = \alpha_4(\alpha_5(x, a_2), a_3)$.

It is easy to prove that

$$f \circ h_1 = T_1, g \circ h_2 = T_3, f_2 \circ g_1 = T_3$$
 and $h \circ f_1 = T_1$

Taking into account the above results we have

$$\begin{array}{ll} \alpha_1(x,z) = f(x) + g(z), & \alpha_2(x,y) = f^{-1}(T_2(y) + T_1(x)) \\ \alpha_3(z,u) = g^{-1}(T_3(z) + T_4(u)), & \alpha_4(z,u) = f_2(z) + T_4(u) \\ \alpha_5(x,y) = f_2^{-1}(T_2(y) + h(x)), & \alpha_6(x,z) = h^{-1}(T_1(x) + T_3(z)). \end{array}$$

The converse is clear.

References

- [1] Petrescu A., G-n-quasigroups, ICTAMI 2007.
- [2] Petrescu A., G n-quasigroup and functional equations on quasigroups, IC-TAMI 2007.
- [3] Sokats'kyi F.M., On the classification of functional equations on quasigroups, Ukrain. Math. Journal, Vol. 56, No.9 (2004), 1259-1266.